



E-ISSN: 2321-2187

P-ISSN: 2394-0514

[www.florajournal.com](http://www.florajournal.com)

IJHM 2022; 10(3): 26-32

Received: 08-02-2022

Accepted: 20-03-2022

**Abdul Qudoos**

Department of Animal Nutrition  
and Nutritional Diseases,  
Institute of Health Science,  
Faculty of Veterinary Medicine,  
Afyon Kocatepe University,  
Afyonkarahisar, Turkey

**Ismail Bayram**

Department of Animal Nutrition  
and Nutritional Diseases,  
Institute of Health Science,  
Faculty of Veterinary Medicine,  
Afyon Kocatepe University,  
Afyonkarahisar, Turkey

## Effects of *Myrtus* and *Nigella sativa* extracts on performance and metabolism of rats: A review

**Abdul Qudoos and Ismail Bayram**

**DOI:** <https://doi.org/10.22271/flora.2022.v10.i3a.818>

### Abstract

In the contemporary world due to technological development the human life has become vulnerable and more at risk because now the diseases are more complicated. Similarly the causative agents, particularly the viruses, have now become more complex due to mutation of their genetic material that results in more resistant and virulent strains. So to counter these diseases a strong immune system is required which can save the living body. Since medieval times the plants have been the main source of food, shelter, and to treat different diseases as well as enhancing the immunity status of both humans and animals. In every civilization the plant-based medicines or phytomedicine have been an important part of first aid. This review elaborates the importance of phytochemical substances i.e. myrtle plant and *Nigella sativa* by exerting positive effects on metabolism and particularly enhancing the immunity of rats which is used as a human model.

**Keywords:** Immunity, metabolism, phytomedicine, rat

### 1. Introduction

#### 1.1 Significance of phytomedicines

The use of plants as medicine dates back to the ancient European, Asian and African civilizations. The use of herbs to treat different diseases is the oldest form of ethno-medical practice known to the human beings and this way of treatment has been an integral part of healthcare of every civilization because at that time it was the only available way of treatment. Since medieval times the herbs are being used to potentiate the immune system and to treat the diseases of humans and animals. In contemporary world the significance of phytomedicines is increased even more because of high prices, side effects, untoward and detrimental effects of synthetic medicines used in humans as well as in the animals to treat various diseases. It is interesting to know that many modern day drugs i.e. codeine, morphine, atropine, quinine, etc. are plant-based. These phytomedicines are unique in a way that these are organic and do not have any side effects if used within the recommended dose. The synthetic medicines have wide range of detrimental side effects whereas their prolonged and consecutive use can create serious health problems for humans as well as for animals. The phytomedicines can also be used as nutraceuticals, which is defined as the food items used to treat the diseases <sup>[1]</sup>.

The phytomedicine contain chemicals compounds which are used as immunostimulants to potentiate the immune system by enhancing immunity status, by activating cytotoxic T-lymphocytes, and by releasing various cytokines toward antigen to achieve the optimum response <sup>[2]</sup>. It is interesting to know that, particularly in the under develop and developing countries, about 70%-80% people prefer to use plant-based drugs <sup>[3]</sup>. The polyphenols exhibit unique biological activity to trigger the signaling pathway, to initiate and to potentiate the immune system to effectively fight the antigens entering the body <sup>[4]</sup>.

The world has badly been affected by the novel corona virus Covid-19 and killed millions of people throughout the world. The scientists throughout the world made vaccines of different brands against the delta strains of corona virus and effectively saved the people. The corona virus has ability to change its genetic material and make new strain, Omicron, now the scientists are trying to make vaccine which is effective against this strain. There is no treatment effective against the virus whereas supportive therapy and a strong immune system can only save living body against viral infection. In this regard polyphenols were considered effective Covid-19 protease inhibitors in trial where affectivity of medicinal plants was evaluated against Covid-19 Mpro which was performed through the exclusive technique of molecular docking. In the trial it was noted different phyto bioactive compoundse. gapiin, rutin, diacetylcurcumin, and diosmi etc.

**Corresponding Author:****Abdul Qudoos**

Department of Animal Nutrition  
and Nutritional Diseases,  
Institute of Health Science,  
Faculty of Veterinary Medicine,  
Afyon Kocatepe University,  
Afyonkarahisar, Turkey

Effectively inhibited the protease enzyme, so it was concluded that the flavonoids might decrease the symptoms of Covid-19 infection [5].

### 1.2 Mechanism of action of phytomedicine

Generally the mode of action of phytomedicines includes diuretic, antihypertensive, anti-diabetic, immuno potentiating, anti-microbial, gastro protective, antioxidant, anti-inflammatory, etc. However exact mechanism of action is still not very much clear whereas it is supposed that it may be related to chemical composition of compound. It is considered when the phyto bioactive compounds are added to diet of animals they exert beneficial effects on the beneficial gut microbiota, decrease the microbial toxic metabolites, and show antimicrobial effects on the intestinal pathogenic bacteria [6]. In addition to this the phytochemicals also exhibit their mode of action through immunomodulation by increasing immune cells, modulation of cytokines and increasing the production of antibodies [7].

### 2. Reasons for using laboratory animals for research purpose

The rat is the most suitable laboratory animal to be used as a human model, and it is commonly utilized for therapeutic purposes. The animals are used by researchers to study human physiology and health related medical problems, as well as to guarantee the affectivity of novel clinical treatments. A portion of these issues are simply conceivable to research in living bodies and whenever there is no other choice left the scientists use animals. The animals have biological similarity to human beings (the DNA of mice resembles almost over 98% to that of humans) and are susceptible to a large number of similar infections. The rats have short life cycles, permitting them to be examined throughout an extensive stretch of time or through various generations. Besides, the researchers can control animal's environmental factors (diet, temperature, lighting, etc.), which is almost difficult when human are under study. The most fundamental contention for utilizing animals is on the grounds that it would be unethical to intentionally subject the humans to health hazards to study how diseases develop.

### 3. *Nigella sativa*

The plant *Nigella sativa*, also called black seeds, member of Ranunculaceae botanical family and due to its wide range of pharmacological characteristics, historical and religious background it is recognized as a miracle herb. It is mostly cultivated in South Europe, Northern African countries and Southwest Asian countries, Mediterranean region, Syria, Pakistan, India, Turkey and Saudi Arabia [8]. It is an annual flowering plant, 20-30 cm length having linear and leaves are of lanceolate shape. Flowers contain 5-10 petals with variety of colors such as white, yellow, blue, pink. It has fruit that is composed of 3-7 follicles having many black colored, trigonous, oblong, angular seeds which have bitter taste, aromatic smell and dimensions include length 0.2 cm and width 0.1 cm. Both, seed and oil, are famous to be used as traditional medicine whereas *Nigella sativa* has lot of importance in Unani and Ayurvedic traditional and herbal treatment system [9]. The *Nigella sativa* is used for its unique and wide range of biological and therapeutic properties including diuretic, antihypertensive, antidiabetic, immunomodulatory, analgesic, anthelmintic, renal and antioxidant, gastroprotective, anti-inflammatory, bronchodilator, antimicrobial, etc. [10]. Thymoquinone is

important phytoactive chemical compound and responsible for most of therapeutic activities [11].




Fig 1: *Nigella sativa* (A) flower (B) seeds

### 4. Religious importance

The black seeds or black cumin has religious importance as well and according to the Tibb-e-Nabwi (the Prophetic Medicine) it is prescribed to use it regularly to lead a healthy life. In the Islamic scripture of the book of Sahih Al-Bukhari there is a famous hadith by the holy Prophet Muhammad (PBUH) narrated by Hazrat Abu Hurairah (RA), "There is cure in the black seeds for every disease excluding death" [12].

### 5. Chemical composition

*Nigella sativa* contains following proportions of important phytoactive compounds including the most abundantly found thymoquinone TQ (30%-48%, responsible for majority of the pharmacological activities), whereas the other phytochemicals include dithymoquinone, thymohydroquinone, carvacrol (6%-12%), longifolene (1%-8%), p-cymene (7%-15%), t-anethol (1%-4%), 4-terpineol (2%-7%),  $\alpha$ -pinene and thymol, etc. as well as some other important biochemical constituents e.g. limonene, citronellol, carvone, etc. which are mostly present in essential oil and these are responsible for antioxidant potential to ameliorate the oxidative stress [13]. Furthermore insulinotropic characteristics, minimizing the cholesterol level, maintaining the  $\beta$ -cells integrity and neutralizing the untoward effects of drug are also associated with phytoactive compounds of black cumin [14].

### 6. Nutritional and phytochemical composition

*Nigella sativa* is composed of fatty oil (35.6-41.5%), fat (28.5%), protein (26.7%), ash (4.8%), crude fiber (8.4%), carbohydrates (24.9%), and volatile oil (0.5-1.6%), (Ahmad, et al., 2013), moisture (8.1-11.6%) and cellulose (6.8-7.4%) [15]. Vitamins and minerals are present in berries in considerable proportions. The vanillic acid is present in roots whereas carotene is present in seeds. The unsaturated fatty acids include dihomolinoleic acid (10%), oleic acid (20%), eicodadienoic acid (3%), and linolic acid (50-60%) whereas main saturated fatty acids include stearic acid and palmitic acid in which stigmaterol (6.57-20.92%) and  $\alpha$ -sitosterol (44-54%) are pioneers [16]. In addition to this other fatty acids present in black seeds include palmitoleic acid, linolenic acid, arachidonic acid, linoleic acid, campesterol,  $\Delta^5$ -avenasterol,  $\beta$ -sitosterol,  $\Delta^7$ -stigmaterol, and  $\Delta^7$ -avenasterol [15].

The phytochemical analysis, as reported by [17] shows a diverse group of other phytoactive compounds i.e. nigellidine, carvone, nigellone, cycloeucalenol, citrostradienol, lophenol, gramisterol, stigmastanol, ostusifoliol, butyrospermol,  $\beta$ -amyrin, tirucallol, taraxerol,

cycloartenol, 3-O-[ $\beta$ -D-xylopyranosyl (1 $\rightarrow$ 3)- $\alpha$ -L-arabino-pyranosyl]-28-O-[ $\alpha$ -L-rhamnopyranosyl(1 $\rightarrow$ 4)- $\beta$ -D-glucopyranosyl(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranosyl] hederagenin,  $\beta$ -unsaturated hydroxyl ketone, esters of dehydrostearic and linoleic acid, melanthigenin, melanthin, resin, 3-O-[ $\beta$ -D-xylopyranosyl (1 $\rightarrow$ 2)- $\alpha$ -L-rhamnopyranosyl (1 $\rightarrow$ 2)- $\beta$ -D-glucopyranosyl]-11-methoxy-16, glycosidalsaponin, stigma-5, 22-dien-3- $\beta$ -D-glucopyranoside, nigellidine-4-O-sulfite, 23-dihydroxy-28-methylolean-12-enoate, cycloart-23-methyl-7, 20, 22-triene-3 $\beta$ , 25-diol, N. mines A1, a2, B1, B2, N. mines A3, a4, A5, and C [16], octahydropyrazino [2, 1-a:5, 4-a'] diisoquinoline, decanynigelloic acid diglucoside [n-decanyl-3-aldehydic-4-methoxy-5-hydroxy benzoate-5- $\beta$ -D-glucopyranosyl (2 $\rightarrow$ 1)- $\beta$ -D-glucopyranosyl(2 $\rightarrow$ 1)- $\beta$ -D-glucopyranoside]] and nigelabdienoyltriglucoide [homolabd-5, 9(11)-dien-16-onyl- $\beta$ -D-glucopyranosyl (2 $\rightarrow$ 1)- $\beta$ -D-glucopyranosyl (2 $\rightarrow$ 1)- $\beta$ -D-glucopyranoside] [18].

## 7. Pharmacological effects

### 7.1. Antibacterial Effect

A trial was performed in rats where the extracts of *Nigella sativa*, together with streptomycin and gentamicin, showed synergistic effect and successfully minimized the infection by *staphylococcus* spp. In the trial filter paper discs, having extract of black cumin @ 25-400  $\mu$ g/disc, were used to exhibit antibacterial characteristics and resultantly the concentration of Gram positive bacteria (*Staphylococcus aureus*) was reduced [19].

### 7.2. Antiviral effects

*Nigella sativa* extract was reported to increase the T4, T8 helper cells, and also increase the cellular activity by natural killer cell [20]. The mice was experimentally infected with murine cytomegalovirus and in order to treat the black cumin oil was intraperitoneally administered which improved the immune system by increasing the T lymphocytes and INF- $\gamma$  production [21].

### 7.3. Antioxidant effects

Herbal supplements' antioxidant characteristic prevent the living bodies against fatal diseases including cancer, cardiovascular and neurodegenerative diseases. A research trial was conducted where the phytoactive compounds (thymoquinone, carvacrol, tanethole, and 4-terpineol) of black cumin enhanced the antioxidant status and exhibited the scavenging characteristics [22].

### 7.4. Anti-diabetic Effect

In carbohydrate and lipid metabolism, efficacy of L-carnitine and  $\alpha$ -lipoic acid ( $\alpha$ -LA), along with black seed, was investigated in rats which were administered streptozotocin @ 65 mg/kg intraperitoneally. The glucose metabolism can be determined by the blood glucose level in fasting situation, insulin, C-peptide, and the activity of pyruvate dehydrogenase enzyme. In this situation  $\alpha$ -LA and the black cumin both have potential to minimize the elevated blood glucose level. In diabetic rats, a considerable improvement is seen in the carbohydrate metabolism when the rats were offered mixture of  $\alpha$ -LA, L-carnitine and black seeds.

The diabetes was reported to increase concentrations of malondialdehyde MDA, glucose and decrease insulin's proportion and superoxide dismutase. To counter this condition, the aqueous extract and oil of black cumin both were administered and the effects were evaluated on the serum insulin, glucose levels, SOD and MDA level in

pancreatic tissues of streptozotocin-induced diabetic rats. It was reported that the treatment successfully minimized the diabetes-induced increase of MDA, and remarkably increased the insulin and superoxide dismutase levels. The thymoquinone was reported to affect at molecular level by neutralizing streptozotocin detrimental effects e.g. heterochromatin material, mitochondrial vacuolization and fragmentation. The results showed that black cumin extract and thymoquinone imparted the anti-diabetic impacts diabetic rats by decreasing the oxidation stress along with maintaining the integrity of pancreatic  $\beta$ -cells [23].

## 8. Myrtle (*Myrtus communis* Linn)

This plant is a spontaneous, evergreen and an important member of Mirtaceae and subfamily Myrtoideae which contains 145 genera and almost 5500 species and mostly grows in Mediterranean region. The genus of this family is composed of flowering plant having 16 species which are found in Middle East and Asian countries. The rosemary residues, thyme residues and their EO are widely used in the feed of animals [24] but that of myrtle has not been widely reported [25].

Myrtle is an evergreen sclerophyll small tree with minimum length of 1.8-2.4 centimeter [26]. There are thick, ovate or lanceolate green 3.5 cm long leaves which contain important phytochemicals such as tannins, flavonoids and EO [27]. The fruit of myrtle can be of two specific colors - dark or white, and most of the times the fruits are of dark color [28]. The dietary usage of myrtle plant in animals may be useful for extending the shelf life and also for maintaining the meat and carcass quality.



Fig 2: Myrtle leaves and berries

## 9. Chemical composition

Myrtle plant's leaves are composed of linalool (7.7-15.8%), 1, 8-cineole (13.5-19.6%), terpinolene, linalyl acetate (2.5-6%), flavonoid and tannins [29]. The fruits contain fatty acids organic acids (9-52%) and anthocyanins (0.2-54%) [30]. In myrtle  $\alpha$ -pinene (~10-60%) and 1, 8-cineole (~12-34%) compounds are commonly found in stem, flowers and leaves [31].

The essential oils are found in hairs, glands, cavities, resin ducts or ducts [32]. Although the essential oils are important phytochemicals but generally their concentration in total mass of the plant rarely exceeds to 1% [33]. The proportion of EO in stem, flower and leaf is 0.08%, 0.30% and 0.61% respectively [31]. The EO does not have any color, or slight yellow, hydrophobic, non-polar, less dense as compared to water, and soluble in alcohol [34]. The oils are composed of Z-3-hexenol, E-2-hexenal,  $\alpha$ -thujene, hexanol, tricyclene,  $\alpha$ -pinene,  $\beta$ -pinene, sabinene,  $\alpha$ -terpinene,  $\delta$ -3-carene, myrcene, p-cymene, E- $\beta$ -ocimene, 1,8-cineole, limonene, terpinolene, E-oxide, borneol, linalool, terpinene-4-ol, p-cymene-8-ol, nerol,  $\alpha$ -terpineol, myrtenol, cis-carveol, bornyl acetate, geraniol,



eugenol, myrtenyl acetate, linalyl acetate, geranyl acetate, methyl eugenol,  $\alpha$ -terpinyl acetate,  $\beta$ -caryophyllene, neryl acetate, germacrene-D, alloaromadendrene,  $\alpha$ -humulene, spathulenol, thiophene, nonadecane,  $\beta$ -elemene, geranyl 2-methylbutyrate, camphene,  $\alpha$ -phellandrene, cis-linalool oxide, caryophylleneoxide, tridecane and trans-linalool oxide. Mainly the EO is divided into below mentioned categories [35-36].

- Terpenes (sesquiterpene hydrocarbons and monoterpene hydrocarbons)
- Terpenoids (oxygenated sesquiterpenes and oxygenated monoterpenes)
- Phenylpropanoids

Similarly *Myrtus communis* plant extracts are composed of polyphenols, and these chemical compounds are divided into 3 main categories.

- Phenolic acids include ellagic, garlic, vanillic, ferulic acid and syringic)
- Tannins include condensed tannins (Proanthocyanidins) and hydrolysable tannins (gallotannins)
- Flavonoids include quercetin, myricetin, catechin, and their respective derivatives which include myricetin-3-d-galactoside, myricetin-3-d-rahmnoside, quercetin-3-rutinoside, catechin derivatives (Flavanols) and quercetin-3-d-rahmnoside [31, 37].

## 10. Pharmacological effects

### 10.1. Antibacterial effect

In a trial the myrtle crude extracts decreased the growth of *Micrococcus luteus*, *Staphylococcus aureus*, *Streptococcus pyogenes*, *Proteus vulgaris*, *Streptococcus agalactiae*, *E. coli*, *Listeria monocytogenes*, and *Campylobacter jejuni*. Minimum inhibitory concentration (MIC) ranged from 0.1 for *S. aureus* and *M. luteus* to more than 2 mg/ml for *E. coli*. The myrtle extracts exhibited increased MIC of 0.025 mg/ml for *Staphylococcus aureus* and *Micrococcus luteus* and MIC of 0.1 mg/ml for *Escherichia coli*. The myrtle essential oils exhibited resistance activity against *M. tuberculosis* but did not show any activity for *M. avium* [38]. An enhanced antibacterial activity can be exhibited during synergistic effects of essential oils and bacteriophages. The combined impact resulted in damaging the cell wall of bacteria by EO either penetrates the bacteriophage in cells or stimulatory

effects of bacteriophage and essential oil on cell membranes [39].

### 10.2. Virucidal activity

Essential oils also contain an important characteristic of destroying or dismantling the viruses by denaturing the viral glycoproteins [40]. According to [41] there are below mentioned two possibilities of mode of action.

- Inhibition of specific processes in the replication of virus.
- Masking viral components using which the virus adsorbs and penetrates into the host cell.

In addition to the above mentioned possible modes of action, some phytochemicals also function by inhibiting the synthesis of nucleic acid or inhibiting stages in virus's multiplication process [42]. The plants of this family contain an important antiviral compound  $\alpha$ -caryophyllene [40].

### 10.3. Gastrointestinal effects

A research trial was performed by [43] where black seed extract was given to the laparotomized Wistar rats @ 25, 50, and 100 mg/kg, and the findings revealed below mentioned results.

- Healed the esophageal damage.
- Reduced malondialdehyde MDA concentration.
- Potentiated some enzymes e.g. CAT, SOD, etc.
- Increased the plasma scavenging activities.
- Involved in regulating the calcium ions' balance ultimately balancing and restoring the ion homeostasis.

Similarly the Wistar rats were experimentally given ethanol or indomethacin and an aqueous extract of myrtle barriers was given to the rats which reduced the ulcer index, and gastric acidity [44]. In a similar research trial the Wistar rats were experimentally given castor oil and then given myrtle aqueous extract @ 25, 50 and 100 mg/kg which reduced diarrhea, minimized the oxidative stress, and potentiated the antioxidative enzymatic activities [43]. In another trial on the Wistar rats, juice of myrtle barriers was used @ 5 and 10 ml/kg which reduced gastric emptying, gut activity, prevented diarrhea, lowered stress of oxidative process, increased the enzymatic activities, etc. [46].

**Table 1:** Effects of extract of *Myrtus communis*

Extract	Wistar rat model	Dose rate	Impacts	Reference
Aqueous extract	Laparotomized male	25, 50, and 100 mg/kg for 6 hours	Healing of histopathology Low oxidation stress Enhanced antioxidative enzymatic activity Reinstatement of homeostasis of ions	[43]
Aqueous and methanolic extract of berries	Gastric ulcer induced by ethanol	105 and 175 mg/kg 93 and 154 mg/kg for 1-5 hour	Reduced ulcer, mucus, acidity Healing of the histopathology	[44]
Juice of berry	Diarrhea induced by castor oil	5-10 ml/kg for 1 hour	Lower the chances of diarrhea Lower the oxidation stress Enhanced activity of enzymes	[46]

### 10.4. Anti-diabetic effects

The phytoactive compounds are considered very effective in lowering or preventing the risk of diabetes mellitus which is an important disease regarding the human health [47]. In a trial aqueous extract of black cumin was used @ 400 and 800 mg/kg in rats and the results showed significant effects on levels of glucose, urea, serum lipid and creatinine level, and liver enzyme [48]. In another trial the myrtle essential oil was used in alloxan-induced diabetic rabbits and findings of trial

revealed significant reduction in glucose level, triglyceride level, hepatic nitrite, and also affected glucokinase and concentration of glycogen whereas the serum insulin level was not affected [49].

The black cumin extract have potency to reduce the chances of diabetes mellitus in streptozotocin-induced diabetic rats whereas the extract didn't show its effects on blood glucose level when used on normal rats. Furthermore EO of myrtle leaves can decrease glucose level in type-2 diabetes and can


induce hypoglycaemic (by enhancing glycolysis, glycogenesis and by decreasing glycogenolysis) and mild hypotriglyceridemic condition in diabetic animals. The myrtle essential oil has the ability to decrease the intestinal glucose uptake as well as inhibit  $\alpha$ -glycosidase enzyme which has the potency to induce hypoglycemia in diabetic rabbits [50]. Myrtle leave extract @ 800 mg/kg BW was used on diabetic rats and the results revealed antihyperglycaemic effects, moreover the administration also had significant effects on blood biochemical parameters such as triglycerides, cholesterol, glucose, HDL, LDL, AST, ALT, BUN, total proteins, creatinine, albumin and globulin [51].

### 10.5. Anti-inflammatory effects

The myrtle plant contained anti-inflammatory activity as it

was reported in a research trial where the myrtle plant extracts @ 10 ml/kg was used in rats [52]. Similarly black cumin extract protected the albino mice against xylene-induced inflammatory response [53]. The myrtle seeds @ 25, 50, and 100 mg/kg was used on ethanol intoxicated rats which descended the concentration of IL-8, TNF- $\alpha$ , IL-1 $\beta$ , IL-6, and lipid peroxidation, etc. [54].

In an in-vivo and in-vitro trial on alloxan induced rats, the decoction of Sahara myrtle was orally administered @ 100 and 300 mg/kg BW, along with Glibenclamide @ 5 mg/kg (reference drug), to check its effects and the results revealed significant antihyperglycaemic effects at high dose rates. It was concluded that the diabetes lowering activity of Sahara myrtle was possibly because of high phenolic, tannin, and flavonoid contents [55].



**Fig 3:** The potential positive effects of myrtle on the health of rat

## 11. References

1. Bjelakovic G, Nikolova D, Gluud C. Antioxidant supplements and mortality. *Current Opinion in Clinical Nutrition and Metabolic Care*. 2014;17(1):40-44.
2. Saxena M, Saxena J, Nema R, Singh D, Gupta A. Phytochemistry of medicinal plants. *Journal of pharmacognosy and phytochemistry*, 2013; 1(6).
3. Hamilton AC. Medicinal plants, conservation and livelihoods. *Biodiversity & Conservation*. 2004;13(8):1477-1517.
4. Ding S, Jiang H, Fang J. Regulation of Immune Function by Polyphenols. *Journal of Immunology Research*, 2018, 1-8. DOI:10.1155/2018/1264074.
5. Bhaliya J, Shah V. Identification of Potent COVID-19 Main Protease (Mpro) Inhibitors from Curcumin Analogues by Molecular Docking Analysis. *Int. J. Adv. Res. Ideas Innov. Technol*. 2020;6(2):664-672.
6. Kim JE, Lillehoj HS, Hong YH, Kim GB, Lee SH, Lillehoj EP, *et al*. Dietary Capsicum and *Curcuma longa* oleoresins increase intestinal microbiome and necrotic enteritis in three commercial broiler breeds. *Research in Veterinary Science*. 2015;102:150-158. DOI:10.1016/j.rvsc.2015.07.022.
7. Kim DK, Lillehoj HS, Lee SH, Jang SI, Park MS, Min W, *et al*. Immune effects of dietary anethole on *Eimeria acervulina* infection. *Poultry Science*. 2013;92(10):2625-2634. DOI:10.3382/ps.2013-03092.
8. Khare CP. Encyclopedia of Indian medicinal plants: rational western therapy, ayurvedic and other traditional usage, botany. Springer, 2004.
9. Goreja WG. Black seed. *Nature's Miracle, Remedy Amazing Herbs Press*, New York, 2003, 1-64.
10. Abel-Salam BKA. Immunomodulatory effects of black seeds and garlic on alloxan-induced diabetes in albino rat. *Allergologia et immuno pathologia*. 2012;40(6):336-340.
11. Al-Ali A, Alkhawajah AA, Randhawa MA, Shaikh NA. Oral and intraperitoneal LD50 of thymoquinone, an active principle of *Nigella sativa* in mice and rats. *J Ayub Med Coll Abbottabad*. 2008;20(2):25-27.
12. Sahih Al-Bukhari. *Book of Medicine*. Chapter No. 7, Hadith No. 5688. Darussalam Publishers and Distributors. Kingdom of Saudi Arabia. 1997;7:330.
13. Butt MS, Sultan MT. *Nigella sativa* reduces the risk of various maladies. *Crit Rev Food Sci Nutr*. 2010;50:654-665.
14. Alenzi FQ, El-Bolkiny YS, Salem ML. Protective effects of *Nigella sativa* oil and thymoquinone against toxicity induced by the anticancer drug cyclophosphamide. *Br J Biomed Sci*. 2010;67:20-28.
15. Gharby S, Harhar H, Guillaume D, Roudani A, Boulbaroud S, Ibrahim M, Charrouf. Chemical investigation of *Nigella sativa* L. seed oil produced in Morocco. *Journal of the Saudi Society of Agricultural Sciences*. 2015;14(2):172-177.
16. Ahmad A, Husain A, Mujeeb M, Khan SA, Najmi AK, Siddique NA, *et al*. A review on therapeutic potential of *Nigella sativa*: A miracle herb. *Asian Pacific journal of tropical biomedicine*. 2013;3(5):337-352.

17. Karna SKL. Phytochemical screening and gas chromatography-mass spectrometry and analysis of seed extract of *Nigella sativa*, Linn. Int J Chem Stud. 2013;1(4):183-187.
18. Ali B, Ali M, Amin S, Mir SR. Enhancement of gut permeation of amoxicillin with *Nigella sativa* seed extract and its phytochemical screening. Chinese journal of natural medicines. 2018;16(2):125-130.
19. Hanafy MSM, Hatem ME. Studies on the antimicrobial activity of *Nigella sativa* seed (black cumin). Journal of ethnopharmacology. 1991;34(2-3):275-278.
20. El-Kadi A, Kandil O. Effect of *Nigella sativa* (the black seed) on immunity. In Proceeding of the 4<sup>th</sup> International Conference on Islamic Medicine, Kuwait. Bull Islamic Med. 1986;4:344-8.
21. Salem ML, Hossain MS. Protective effect of black seed oil from *Nigella sativa* against murine cytomegalovirus infection. International journal of immunopharmacology. 2000;22(9):729-740.
22. Burits M, Bucar F. Antioxidant activity of *Nigella sativa* essential oil. Phytotherapy research. 2000;14(5):323-328.
23. Salama RHM. Hypoglycemic effect of lipoic acid, carnitine and *Nigella sativa* in diabetic rat model. International journal of health sciences. 2011;5(2):126-134.
24. Smeti S, Hajji M, Mekki I, Mahouachi M, Atti N. Effects of dose and administration form of rosemary essential oils on meat quality and fatty acid profile of lamb. Small Ruminant Research. 2018;158:62-68. DOI:10.1016/j.smallrumres.2017.10.007
25. Tibaoui S, Essid I, Smeti S, Bertolin JR, Joy M Atti, N. Fatty acid profile, physiochemical properties and oxidative stability of ewe's sausage as affected by distilled myrtle (*Myrtus communis*) leaves' intake. International Journal of Food Science Technology. 2020;55(3):1151-1161. DOI:10.1111/ijfs.14549
26. Mendes MM, Gazarini LC, Rodrigues ML. Acclimation of *Myrtus communis* to contrasting Mediterranean light environments effects on structure and chemical composition of foliage and plant water relations. Environmental and Experimental Botany. 2001;45(2):165-178.
27. Baytop T. Therapy with medicinal plants in Turkey (past and present). Publication of the Istanbul University, 1999.
28. Klein JD, Cohen S, Hebbe Y. Seasonal variation in rooting ability of myrtle (*Myrtus communis* L.) cuttings. Scientia Horticulturæ. 2000;83(1):71-76.
29. Gardeli C, Vassiliki P, Athanasios M, Kibouris T, Komaitis M. Essential oil composition of *Pistacia lentiscus* L. and *Myrtus Communis* L.: Evaluation of antioxidant capacity of methanolic extracts. Food Chemistry. 2008;107:1120-1130.
30. Messaoud C, Laabidi A, Boussaid M. *Myrtus communis* L. infusions: the effect of infusion time on phytochemical composition, antioxidant, and antimicrobial activities. Journal of Food Science. 2012;77(9):C941-C947. DOI:10.1111/j.1750-3841.2012.02849.x
31. Wannes WA, Mhamdi B, Sriti J, Jemia MB, Ouchikh O, Hamdaou G, et al. Antioxidant activities of the essential oils and methanol extracts from myrtle (*Myrtus communis* var. *italica* L.) leaf, stem and flower. Food and chemical toxicology. 2010;48(5):1362-1370. DOI:10.1016/j.fct.2010.03.002
32. Liolios CC, Graikou K, Skaltsa E, Chinou I. Dittany of Crete: a botanical and ethnopharmacological. Journal of Ethnopharmacology. 2010;131(2):229-241.
33. Bowles EJ. The chemistry of aroma therapeutic oils. 3<sup>rd</sup> ed. South Australia: Griffin Press, 2003.
34. Gupta V, Mittal P, Bansal P, Khokra SL, Kaushik D. Pharmacological Potential of *Matricaria recutita*. Int J Pharm Sci Drug Res. 2010;2(1):12-16.
35. Andrade EHA, Alves CN, Guimaraes EF, Carreira LMM, Maia JGS. Variability in essential oil composition of *Piper dilatatum* L. C Rich. Biochemical Systematics and Ecology. 2011;39(4-6):669-675. DOI:10.1016/j.bse.2011.05.021
36. De Sousa DP. Analgesic-like activity of essential oils constituents. Molecules. 2011;16(3):2233-2252. DOI:10.3390/molecules16032233
37. Tuberoso CIG, Rosa A, Bifulco E, Melis MP, Atzeri A, Pirisi FM, et al. Chemical composition and antioxidant activities of *Myrtus communis* L. Berries extracts. Food Chemistry. 2010;123(4):1242-1251. DOI:10.1016/j.foodchem.2010.05.094
38. Zanetti S, Cannas S, Molicotti P, Bua A, Cubeddu M, Porcedda S, et al. Evaluation of the antimicrobial properties of the essential oil of *Myrtus communis* L. against clinical strains of *Mycobacterium* spp. Interdisciplinary Perspective of Infectious Diseases, 2010, 1-4. DOI:10.1155/2010/931530
39. Kon KV, Rai MK. Plant essential oils and their constituents in coping with multidrug-resistant bacteria. Expert review of anti-infective therapy. 2012;10(7):775-790.
40. Djilani A, Dicko A. The therapeutic benefits of essential oils. Nutrition, well-being and health. 2012;7:155-179.
41. Saddi M, Sanna A, Cottiglia F. Antiherpesvirus activity of *Artemisia arborescens* essential oil and inhibition of lateral diffusion in Vero cells. Annals of Clinical Microbiology and Antimicrobials. 2007;6:10. DOI:10.1186/1476-0711-6-10
42. Jassim SAA, Naji MA. Novel antiviral agents: a medicinal plant perspective. Journal of Applied Microbiology, 2003; 95(3), 412-427. doi:10.1046/j.1365-2672.2003.02026.
43. Jabri MA, Rtibi K, Sakly M, Marzouki L, Sebai H. Role of gastrointestinal motility inhibition and antioxidant properties of myrtle berries (*Myrtus communis* L.) juice in diarrhea treatment. Biomedicine & pharmacotherapy. 2016;84:1937-1944.
44. Sumbul S, Ahmad MA, Asif M, Saud I, Akhtar M. Evaluation of *Myrtus communis* Linn. berries (common myrtle) in experimental ulcer models in rats. Human & experimental toxicology. 2010;29(11):935-944.
45. Md. Shahab Uddin, Nuri ZN. Therapeutic use of *Nigella sativa*: A review. Int. J Horti Food Sci. 2021;3(1):06-12
46. Jabri MA, Rtibi K, Tounsi H, Hosni K, Marzouki L, Sakly M, et al. Fatty acid composition and mechanisms of the protective effects of myrtle berry seed aqueous extract in alcohol-induced peptic ulcer in rat. Canadian journal of physiology and pharmacology. 2017;95(5):510-521.
47. Olfert, MD, Wattick RA. Vegetarian diets and the risk of diabetes. Current diabetes reports. 2018;18(11):1-6.
48. Benkhayal FA, Musbah E, Ramesh S, Dhayabaran D. Biochemical studies on the effect of phenolic compounds extracted from *Myrtus communis* in diabetic rats. Tamil Nadu Journal of Veterinary and Animal Sciences. 2009;5(3):87-93.

49. Sepici-Dincel A, Açıkgoz Ş, Çevik C, Sengelen M, Yeşilada E. Effects of *in vivo* antioxidant enzyme activities of myrtle oil in normoglycaemic and alloxan diabetic rabbits. *Journal of ethnopharmacology*. 2007;110(3):498-503.
50. Sepici A, Gurbuz I, Cevik C, Yesilada E. Hypoglycaemic effects of myrtle oil in normal and alloxan-diabetic rabbits, *J. Ethnopharmacol*. 2004;93(2-3):311-318.
51. Fahim AB, El-Ghaithi M, Amesh S, Dhayabaran D. Biochemical studies on the effect of phenolic compounds extracted from *Myrtus communis* in diabetic rats, *Tamilnadu J Vet Anim Sci*. 2009;5(3):87-93.
52. Al-Hindawi MK, Al-Deen IH, Nabi MH, Ismail MA. Anti-inflammatory activity of some Iraqi plants using intact rats. *Journal of Ethnopharmacology*. 1989;26(2):163-168.
53. Hosseinzadeh H, Khoshdel M, Ghorbani M. Antinociceptive, anti-inflammatory effects and acute toxicity of aqueous and ethanolic extracts of *Myrtus communis* L. aerial parts in mice. *Journal of acupuncture and meridian studies*, 2011;4(4):242-247.
54. Jabri MA, Marzouki L, Sebai H. Myrtle berries seeds aqueous extract abrogates chronic alcohol consumption-induced erythrocytes osmotic stability disturbance, hematological and biochemical toxicity. *Lipids in health and disease*. 2018;17(1):1-10.
55. Farah R, Amira S, el Houda HN, Mahfoud HM. Blood biochemical parameters effect of Sahara myrtle on diabetic rats. *Pharmacophore*. 2019;10(1):71-77.