International Journal of Herbal Medicine
Available online at www.florajournal.com

IJHM
International Journal of Herbal Medicine

E-ISSN: 2321-2187
P-ISSN: 2394-0514
www.florajournal.com
IJHM 2024; 12(6): 126-132
Received: 25-09-2024
Accepted: 28-10-2024

Prakash Patil
Department of Pharmacognosy, Pune City College of Pharmacy, Pune, Maharashtra, India

Corresponding Author:
Prakash Patil
Department of Pharmacognosy, Pune City College of Pharmacy, Pune, Maharashtra, India

Phytochemical and ethnomedicinal significance of *Helicteres isora* in herbal therapeutics

Prakash Patil

DOI: https://doi.org/10.22271/flora.2024.v12.i6b.1024

Abstract
Helicteres isora L. (Malvaceae), commonly known as East Indian screw tree, is a widely used medicinal plant in South and Southeast Asia. Traditional systems—including Ayurveda, Siddha, Unani, and various folk traditions—employ its fruits, roots, stem bark, and leaves to manage gastrointestinal disturbances, metabolic disorders, infections, respiratory ailments, and gynecological conditions. This narrative review synthesizes botanical and ethnopharmacological knowledge with contemporary phytochemical and pharmacological evidence to clarify how *H. isora* contributes to herbal therapeutics. Morphologically, the species is identified by its characteristic twisted, screw-like follicles and pubescent leaves; it is adapted to dry deciduous habitats and often used as a multipurpose species in rural health care and livelihoods. Phytochemically, *H. isora* contains a diverse repertoire of secondary metabolites—principally polyphenols (flavanols, flavonols, tannins), phenolic acids, triterpenoids (e.g., betulinic and oleanolic acid derivatives), steroids, lignans, simple and oligomeric sugars, and alkaloid traces—supported by chromatographic and spectrometric profiling in multiple studies. Across preclinical models, extracts and isolated constituents have demonstrated antioxidant, antimicrobial (bacterial and fungal), antidiarrheal, antidiabetic/α-glucosidase inhibitory, anti-inflammatory, analgesic, hepatoprotective, gastroprotective, and antispasmodic properties. Preliminary safety assessments suggest a reasonable margin of safety for traditional oral doses, although standardized toxicology, herb-drug interaction data, and reproductive toxicity studies remain limited. Quality control considerations—macroscopic/microscopic authentication, marker-based HPTLC/HPLC fingerprints, and contamination limits—are essential due to high use and market adulteration risk. Conservation and sustainable supply are salient because local wild collection pressures threaten populations in certain ranges. Overall, *H. isora* represents a promising candidate for evidence-led, standardized phytomedicine development, especially for gastrointestinal and metabolic indications, provided that future research prioritizes chemotaxonomic clarity, robust bioassay-guided isolation, pharmacokinetics, dose-ranging safety, and multicentric clinical trials.

Keywords: *Helicteres isora*, Malvaceae, ethnomedicine, phytochemistry, polyphenols, triterpenoids, antidiarrheal, antidiabetic, antimicrobial, Ayurveda

1. Introduction
Medicinal plants form the backbone of primary health care for large populations across Asia. Among these, *Helicteres isora* L., known for its distinctive screw-like fruit, enjoys long-standing use in Indigenous and classical medical traditions. In Ayurveda, the plant is referenced for disorders of the gastrointestinal tract (e.g., atisāra/diarrhea, grahaṇī/malabsorption), metabolic imbalance (madhumeha/diabetes), and as a general "stambhana" (astringent) drug due to its tannin-rich profile. Folk medicine in India, Sri Lanka, Nepal, Myanmar, and Thailand similarly employs the plant's bark, root, and fruits for dysentery, intestinal colic, helminthiasis, cough, and as a postpartum restorative.

The modern phytomedicine discourse seeks to bridge such traditional claims with chemical and pharmacological evidence. Over the past two decades, phytochemical investigations have reported polyphenols (catechins, epicatechins, quercetin/kaempferol glycosides), phenolic acids (gallic, ellagic, ferulic, caffeic), triterpenoids (betulinic/oleanolic scaffolds), phytosterols (β-sitosterol), and lignans; these chemistries plausibly underpin antidiarrheal, antimicrobial, antioxidant, and metabolic modulatory effects in preclinical assessments.

This paper reviews the botany, ethnomedicinal knowledge, chemistry, and pharmacology of *H. isora*, identifies translational opportunities and constraints, and highlights quality control, safety, and sustainability considerations relevant to its responsible inclusion in contemporary herbal therapeutics.

2. Botanical Profile and Pharmacognosy

Helicteres isora L. is a deciduous shrub or small tree belonging to the family Malvaceae, usually growing to a height of 3-6 meters, though some specimens can extend up to 8 meters. It is widely known as the "East Indian screw tree," a name derived from its characteristic spirally twisted fruit. The species is native to tropical Asia and is distributed across India, Sri Lanka, Nepal, Myanmar, and Thailand. Within India, it is especially common in the dry deciduous forests and scrublands of Madhya Pradesh, Maharashtra, Chhattisgarh, Andhra Pradesh, and the Western Ghats. Ecologically, it is well adapted to sandy-loam soils, tolerates drought, and is generally found between 200 and 1200 meters above sea level. Flowering occurs between July and September, while the fruits mature from October to January.

The plant exhibits several distinctive morphological features. The leaves are ovate to lanceolate, 6-12 cm long, with serrated margins and a pubescent underside. They show a characteristic palmate venation with five to seven prominent nerves radiating from the base. Flowers are axillary, bisexual, and orange to reddish in color, while the fruits form the most important diagnostic trait: long, slender, twisted follicles measuring 5-10 cm in length and containing multiple angular blackish seeds. The bark is greyish-brown, fissured, and about 0.3-0.5 cm thick with a pronounced astringent taste due to high tannin content. Roots are cylindrical, woody, and pale brown with a faint aromatic odor.

Pharmacognostic investigations have provided detailed insight into the microscopic and chemical characteristics of the plant parts. The bark shows stratified cork and secondary phloem with sclereids and tannin-filled parenchyma. Powder microscopy reveals lignified fibers, stone cells, tannin-rich cells, and calcium oxalate crystals. Leaf powder exhibits unicellular trichomes, polygonal epidermal cells, and anomocytic stomata, while fruits display sclerenchyma tissues and lignified pericarp. These features are crucial in distinguishing authentic raw drug samples from adulterants in herbal markets.

The Ayurvedic Pharmacopoeia of India (2007) has provided official pharmacopoeial standards for *H. isora* fruit, including loss on drying below 12%, total ash not exceeding 10%, and alcohol-soluble extractives above 8%. Tannin content of the fruit is reported to be in the range of 15-20%, while bark tannins range between 12-18%. Marker compounds such as catechin, epicatechin, gallic acid, ellagic acid, betulinic acid, and β-sitosterol have been identified through chromatographic studies and are now considered essential parameters for standardization.

To consolidate these findings, a comparative pharmacognostic profile of different plant parts is presented below.

Table 1: Comparative Pharmacognostic Data of *Helicteres isora*

Plant Part	Macroscopic Features	Microscopic / Powder Characteristics	Major Phytochemical Markers	Standard Values / Data
Leaf	Ovate-lanceolate, 6-12 cm, serrated margins, pubescent underside, palmate venation with 5-7 nerves	Unicellular trichomes, polygonal epidermal cells, anomocytic stomata	Flavonoids (quercetin, kaempferol glycosides), phenolic acids	Moisture content < 12% (API, 2007)
Fruit	Spirally twisted follicle, 5-10 cm, dark brown when mature, astringent taste	Lignified pericarp, sclerenchyma fibers, calcium oxalate crystals	Catechin, epicatechin, gallic acid, ellagic acid, tannins (15-20%)	Alcohol-soluble extractives > 8%; Total ash < 10%
Bark	Greyish-brown, fissured, 0.3-0.5 cm thick, strongly astringent	Stratified cork, secondary phloem with sclereids, tannin-rich parenchyma	Betulinic acid, oleanolic acid, β-sitosterol	Tannin content 12-18% (Gupta *et al*., 2013)
Root	Cylindrical, woody, pale brown, faintly aromatic	Fibers with lignified walls, starch grains, stone cells	Saponin-like fractions, phenolic acids, sterols	Moderate seed germination (40-50%); improved with hot water pre-treatment (ICFRE, 2016)

Trade records further highlight the ethnopharmacological and economic significance of the species. The National Medicinal Plant Board (NMPB) has identified *H. isora* as a high-demand species, with an annual market volume estimated at 200-250 metric tons, primarily of fruits sourced from Madhya Pradesh and Chhattisgarh. The Foundation for Revitalization of Local Health Traditions (FRLHT) also includes *H. isora* among the top 150 medicinal plants in Indian trade. However, increased demand has led to overharvesting and localized population decline. Germination studies conducted by the Indian Council of Forestry Research and Education (ICFRE) demonstrate that the seeds exhibit only 40-50% natural germination, though this can be improved by hot water pre-treatment, underlining the importance of cultivation initiatives for sustainable supply.

Thus, *Helicteres isora* presents a unique pharmacognostic identity, characterized by its morphology, microscopic features, and phytochemical markers. Its recognition in the Ayurvedic Pharmacopoeia, coupled with its significant trade volume and conservation concerns, emphasizes the need for standardized cultivation practices, rigorous authentication, and quality assurance to safeguard its role in herbal therapeutics.

3. Ethnomedicinal Uses

The ethnomedicinal importance of *Helicteres isora* has been recognized for centuries across South and Southeast Asia, where it continues to serve as a primary remedy in both classical and folk systems of medicine. In Ayurveda, the plant is known as *Avartani* or *Avartaki* and is categorized among astringent and digestive drugs, often prescribed for gastrointestinal complaints such as *atisāra* (diarrhea), *grahaṇī* (malabsorption), and *pravāhika* (dysentery). Fruits, when prepared as decoctions, are widely administered for acute diarrhea due to their high tannin content, which helps reduce intestinal secretions and provides a protective effect on the mucosa. In tribal communities of Madhya Pradesh, Chhattisgarh, and Maharashtra, this practice is still prevalent, reflecting its enduring relevance in rural health care.

The uses of *H. isora* extend well beyond the digestive system. Folk healers in Andhra Pradesh and Tamil Nadu prepare powdered fruits with honey to quench excessive thirst and regulate blood sugar, while rural practitioners in Rajasthan and Gujarat use bark decoctions as febrifuges during malarial and seasonal fevers. In Nepal and Sri Lanka, postpartum women are given bark and fruit preparations to regain strength and improve uterine tone, underlining its role in reproductive health traditions. The roots are also valued for

their ability to relieve colic and mild respiratory congestion, often used as infusions in central India.

Topical and pediatric applications further highlight the versatility of the species. Crushed leaves are commonly applied to boils, eruptions, and wounds in Odisha and Jharkhand, while bark poultices are used for joint pain and swelling. In the Western Ghats, fruit pastes are employed against mouth ulcers and throat infections in children. Pediatric use is notable across many tribal belts, where diluted decoctions of fruit are administered to infants for indigestion, flatulence, and worm infestations.

The plant also assumes importance as a community health resource during times of scarcity. In parts of Maharashtra and Madhya Pradesh, fruits are occasionally boiled and consumed as supplementary food during famines, reflecting both its nutritive value and resilience as a survival crop. Such wide-ranging practices underscore the plant's embeddedness in daily healthcare strategies, highlighting its dual role as a therapeutic and cultural resource.

The diversity of ethnomedicinal knowledge is captured in Table 2, which summarizes the regional uses, plant parts employed, modes of preparation, and traditional indications.

Table 2: Ethnomedicinal Uses of *Helicteres isora* Across Regions

Region / Community	Plant Part Used	Mode of Preparation	Condition Treated
Madhya Pradesh, Chhattisgarh (tribal)	Fruit	Decoction	Diarrhea, dysentery
Maharashtra (rural)	Bark	Infusion/decoction	Fever, stomach colic
Andhra Pradesh, Tamil Nadu	Fruit	Powder with honey	Excessive thirst, diabetes
Rajasthan, Gujarat	Bark	Decoction	Malaria, seasonal fevers
Nepal, Sri Lanka	Fruit and bark	Postpartum tonic	Uterine restoration, postnatal weakness
Odisha, Jharkhand	Leaves, bark	Poultice, crushed paste	Skin eruptions, wounds, joint pain
Western Ghats (folk)	Fruit	Paste	Mouth ulcers, throat infection
Central India (tribal pediatric care)	Fruit, bark	Diluted decoction	Flatulence, worm infestation in children
Maharashtra, Madhya Pradesh (scarcity use)	Fruit	Boiled, consumed as food	Famine survival, general weakness

This ethnomedicinal evidence shows that *H. isora* has been used as a multipurpose therapeutic across a broad range of ailments including gastrointestinal, metabolic, febrile, respiratory, dermatological, pediatric, and gynecological conditions. The consistency of these practices across regions and generations indicates a strong empirical foundation. When examined alongside pharmacological studies that confirm its antidiarrheal, antimicrobial, antidiabetic, and anti-inflammatory properties, the traditional uses gain scientific credibility, strengthening the case for developing standardized and evidence-based formulations from this plant.

4. Phytochemistry

The phytochemical composition of *Helicteres isora* has been investigated extensively, revealing a wide range of bioactive secondary metabolites that substantiate its use in ethnomedicine. Early screenings confirmed the presence of tannins, flavonoids, phenolic acids, triterpenoids, steroids, and carbohydrates, while modern chromatographic and spectroscopic analyses have provided detailed identification of individual compounds. These phytochemicals play central roles in the pharmacological activities attributed to the plant, particularly in gastrointestinal disorders, infections, metabolic regulation, and oxidative stress-related conditions.

Polyphenols are among the most abundant constituents. Fruits and bark contain significant levels of catechin and epicatechin, as well as glycosides of quercetin and kaempferol. Ellagitannins and condensed tannins make up 15-20% of the dried fruit, accounting for its astringent taste and its use as an antidiarrheal agent in traditional medicine. Phenolic acids such as gallic acid, ellagic acid, caffeic acid, ferulic acid, and p-coumaric acid are also widely reported and contribute to the antioxidant and antimicrobial properties of the plant.

Triterpenoids and steroids represent another important group. Betulinic acid and oleanolic acid derivatives have been isolated, both with documented anti-inflammatory, hepatoprotective, and gastroprotective potential. Sterols such as β-sitosterol and stigmasterol are present in fruits and bark, further supporting the plant's role in managing metabolic and inflammatory disorders.

Carbohydrates, including glucose, fructose, and sucrose, are present in the fruits, which have occasionally been consumed as a supplementary food during scarcity. Seeds are rich in mucilage, which has soothing effects on the digestive tract. Traces of saponin-like fractions and minor alkaloids have also been detected.

Analytical studies using HPTLC, HPLC, UPLC, and LC-MS/MS have established chemical fingerprints with gallic acid, catechin, and betulinic acid serving as reference markers. Seasonal variation affects metabolite content; fruits harvested in later months of the season often show higher tannin yields. Such findings emphasize the importance of standardization and quality control to ensure therapeutic consistency.

A summary of the major phytoconstituents, their chemical classes, plant parts, and reported biological activities is presented in Table 3.

Table 3: Major Phytoconstituents of *Helicteres isora* and Their Reported Activities

Compound / Group	Chemical Class	Plant Part	Reported Biological Activity
Catechin, Epicatechin	Flavan-3-ols (polyphenols)	Fruit, bark	Antidiarrheal, antioxidant, antimicrobial
Quercetin, Kaempferol glycosides	Flavonols	Leaves, fruits	Anti-inflammatory, antioxidant, antidiabetic
Gallic acid, Ellagic acid, Ferulic acid, Caffeic acid	Phenolic acids	Fruit, bark	Antioxidant, gastroprotective, antimicrobial
Ellagitannins, Condensed tannins	Hydrolyzable/condensed tannins	Fruit	Astringent, antidiarrheal, antimicrobial
Betulinic acid, Oleanolic acid derivatives	Triterpenoids	Bark, fruit	Anti-inflammatory, hepatoprotective, analgesic
β-sitosterol, Stigmasterol	Phytosterols	Bark, fruit	Anti-inflammatory, hypocholesterolemic
Lignans, phenylpropanoids	Lignan-type phenolics	Bark, seeds	Antioxidant, antimicrobial
Glucose, Fructose, Sucrose	Carbohydrates	Fruit	Nutritional value, famine food
Mucilage	Polysaccharide	Seeds	Demulcent, digestive soothing
Saponin-like fractions, trace alkaloids	Secondary metabolites (minor)	Root, bark	Mild antimicrobial, metabolic modulation

The diverse phytochemical repertoire of *H. isora* offers a strong biochemical rationale for its ethnomedicinal applications. Polyphenols and tannins account for its astringent, antimicrobial, and antidiarrheal actions; triterpenoids and sterols support anti-inflammatory and hepatoprotective claims; while mucilage and carbohydrates contribute to digestive benefits and nutritional use. Collectively, these findings demonstrate that the plant's traditional therapeutic roles are closely aligned with its phytochemical architecture, providing a foundation for its potential standardization into evidence-based herbal formulations.

5. Pharmacology and Mechanistic Insights

The pharmacological activities of *Helicteres isora* have been investigated in a variety of experimental systems, revealing a broad spectrum of therapeutic effects that align closely with its ethnomedicinal uses. Extracts prepared from fruits, bark, leaves, and roots have demonstrated antidiarrheal, antimicrobial, antidiabetic, antioxidant, anti-inflammatory, hepatoprotective, gastroprotective, analgesic, and antipyretic properties. These actions can be traced to the synergistic influence of polyphenols, triterpenoids, sterols, and other secondary metabolites.

One of the most well-established traditional uses of *H. isora* is in the treatment of diarrhea and dysentery, and this has been validated through pharmacological studies. Animal models of castor oil-induced diarrhea showed that aqueous and ethanolic fruit extracts significantly reduced fecal output and intestinal motility. The high tannin content is believed to precipitate proteins on the intestinal mucosa, thereby forming a protective layer that decreases secretion and fluid loss. Flavonoids and triterpenoids may also contribute by modulating enteric smooth muscle contractions through calcium channel blockade and inhibition of prostaglandin synthesis, reducing hypermotility and intestinal spasms.

Antimicrobial activity has been reported across several studies. Ethanolic and methanolic extracts exhibited inhibitory effects against *Escherichia coli*, *Shigella dysenteriae*, *Vibrio cholerae*, *Staphylococcus aureus*, and *Candida albicans*. Mechanistic evaluations suggest that phenolic acids and flavonoids disrupt microbial cell walls, interfere with enzyme activity, and generate oxidative stress in pathogens. Minimum inhibitory concentrations (MICs) for ethanolic fruit extracts against enteropathogens have been reported in the range of 250-500 µg/mL, demonstrating practical relevance to their traditional application in infectious diarrhea.

The plant also demonstrates notable antioxidant and anti-inflammatory activity. *In vitro* assays such as DPPH, ABTS, and ferric reducing antioxidant power (FRAP) consistently reveal strong radical scavenging capacities of fruit and bark extracts, attributable to catechin, epicatechin, gallic acid, and ellagic acid. *In vivo* studies have confirmed that treatment with extracts elevates endogenous antioxidant enzymes such as superoxide dismutase, catalase, and glutathione peroxidase while reducing lipid peroxidation markers. These biochemical effects explain its protective role in models of inflammation, where inhibition of pro-inflammatory mediators such as TNF-α, IL-6, COX-2, and LOX pathways has been documented. Such findings validate its traditional use in managing febrile states, body aches, and inflammatory conditions.

Antidiabetic activity has also been highlighted in several preclinical studies. Oral administration of hydroalcoholic fruit extracts improved glucose tolerance in streptozotocin-induced diabetic rats, reduced fasting blood glucose, and decreased glycosylated hemoglobin levels. *In vitro* studies confirmed inhibition of α-glucosidase and α-amylase enzymes, indicating a capacity to delay carbohydrate digestion and reduce postprandial hyperglycemia. Polyphenols and triterpenoids are proposed as the active compounds, with additional antioxidant support protecting pancreatic β-cells from oxidative stress.

Protective effects on the gastrointestinal system have been reported. Ethanolic extracts reduced gastric lesions in ethanol- and NSAID-induced ulcer models, suggesting gastroprotective activity. The mechanism appears to involve enhancement of mucosal defense, increased prostaglandin secretion, and reduction of oxidative damage. Similarly, hepatoprotective effects were observed in toxin-induced liver injury models, where extracts normalized serum enzymes (ALT, AST, ALP) and improved histological features of the liver. These results are attributed to triterpenoids such as betulinic acid and oleanolic acid, which are known to stabilize hepatocyte membranes and counter oxidative stress.

Additional pharmacological actions include analgesic and antipyretic effects demonstrated in rodent models. Extracts prolonged reaction times in hot plate and tail flick tests, and reduced yeast-induced pyrexia, suggesting both central and peripheral mechanisms of pain and fever modulation. Preliminary reports have also indicated mild anthelmintic activity, with seed preparations reducing motility of intestinal worms, though this remains less well studied.

Overall, the pharmacological evidence positions *Helicteres isora* as a multipurpose medicinal plant with a strong correlation between traditional uses and mechanistic insights. The astringent tannins and polyphenols provide antidiarrheal and antimicrobial protection, flavonoids and triterpenoids underpin antioxidant and anti-inflammatory effects, while sterols and phenolic acids contribute to metabolic and gastroprotective benefits. These findings not only affirm its ethnomedicinal reputation but also suggest that the plant holds promise for further development into standardized herbal formulations, provided that more rigorous clinical validation and pharmacokinetic studies are undertaken.

6. Safety, Toxicology, and Interactions

The long history of use of *Helicteres isora* in traditional medicine suggests that the plant has a relatively wide margin of safety at customary doses. Fruits, bark, and roots are routinely used in the form of decoctions, powders, and infusions in rural and Ayurvedic practices, often administered to children, adults, and even postpartum women without reports of acute toxicity. This ethnomedical record provides a foundation for its presumed safety. Nevertheless, systematic toxicological evaluations remain limited, and further research is needed before unequivocal conclusions can be drawn regarding its long-term use.

Preclinical studies available to date indicate low acute toxicity. Ethanolic and aqueous extracts of the fruit and bark have been tested in rodent models, with oral administration up to 2000 mg/kg body weight showing no mortality or major behavioral changes. Subacute administration at moderate doses has not revealed significant alterations in hematological parameters, liver function, or renal function. Histopathological examinations of liver and kidney tissues in these models have also suggested preservation of normal cellular architecture, implying absence of overt toxicity. These findings broadly support its safe use in controlled amounts, although high doses beyond traditional ranges may

carry risks that remain poorly defined.

The pharmacological richness of *H. isora* also implies potential interactions with conventional drugs. The fruit is particularly high in tannins, which are known to form complexes with proteins and minerals. This could reduce the intestinal absorption of certain nutrients such as non-heme iron, and may interfere with the bioavailability of alkaloid-containing medications. Traditional practitioners often administer fruit decoctions after meals to minimize such issues, an observation that reflects indigenous awareness of these interactions. Additionally, *in vitro* studies have shown that extracts of *H. isora* inhibit carbohydrate-digesting enzymes such as α-glucosidase and α-amylase. While this underpins its antidiabetic potential, it also raises the possibility of additive hypoglycemic effects if co-administered with pharmaceutical antidiabetic agents, increasing the risk of hypoglycemia.

Pregnancy and lactation safety remain areas of concern. Although traditional medicine in parts of Nepal and Sri Lanka employs fruit and bark decoctions during the postpartum period, there is insufficient modern toxicological evidence to confirm their safety during pregnancy or breastfeeding. Given the presence of tannins and triterpenoids, caution is advised until systematic reproductive toxicity studies are undertaken. Similarly, data on mutagenicity, genotoxicity, and reproductive outcomes are lacking, which represents an important research gap.

Another important consideration is the potential variability in toxicity and interactions depending on the source, harvest season, and preparation method. Polyphenol and tannin levels, for instance, are known to vary with maturity of the fruit, which could influence both therapeutic efficacy and risk of gastrointestinal irritation when consumed in large amounts. Quality control therefore becomes a vital safety measure, ensuring that standardization of active compounds is achieved and contaminants such as heavy metals, pesticides, or microbial loads are kept within pharmacopeial limits.

7. Herbal Formulations and Dosage Considerations

The therapeutic applications of *Helicteres isora* are not limited to its use as a single drug but extend to its incorporation into compound formulations in Ayurvedic, Siddha, and folk medicine. In the Ayurvedic tradition, the fruits and bark are frequently employed in formulations aimed at treating gastrointestinal disorders such as diarrhea, dysentery, and malabsorption. The powdered fruit (*churna*) is one of the most common dosage forms, administered either alone or in combination with honey, warm water, or ghee to enhance palatability and absorption. Classical texts and ethnomedicinal records indicate a daily dose of 1-3 grams of fruit powder for adults, usually given in divided doses after meals to minimize any potential gastrointestinal irritation. Decoctions (*kvātha*) are also widely prepared by boiling the coarsely crushed fruit or bark in water, generally in the ratio of 1:16, reduced to one-fourth of the volume, with a typical adult dose ranging between 30 and 50 mL twice daily. These preparations are primarily prescribed for diarrhea, dysentery, and febrile conditions associated with intestinal disturbances.

The plant is also present in multi-herbal formulations used for metabolic and respiratory conditions. For instance, in certain Siddha and folk practices of Tamil Nadu, *H. isora* fruit powder is mixed with *Zingiber officinale* and *Piper longum* to alleviate colic and indigestion. In rural Andhra Pradesh, a combination of *H. isora* bark decoction with *Glycyrrhiza glabra* is administered for chronic cough and throat irritation, demonstrating its flexibility in polyherbal prescriptions. Such formulations highlight the synergistic use of *H. isora* with other herbs that share complementary properties.

In pediatric practice, the fruit decoction is commonly diluted and administered in small quantities for indigestion and worm infestations. Traditional healers recommend cautious dosing, often using a half or one-third of the adult dose, given the sensitivity of children to tannin-rich preparations. The use of honey as an adjuvant is frequent, both for improving taste and for enhancing antimicrobial and demulcent activity.

Modern phytomedicine research has suggested the potential for standardized extracts of *H. isora* based on quantifiable marker compounds such as catechin, gallic acid, and betulinic acid. Hydroalcoholic extracts, enriched with polyphenols and triterpenoids, have been explored in preclinical models of diabetes, oxidative stress, and gastrointestinal disorders. However, standardized dosing for such extracts is yet to be clinically established. Extrapolations from preclinical studies suggest that extracts corresponding to 200-400 mg/kg in rodent models may translate into adult human doses of 1.5-3 g/day of crude drug equivalent, though formal dose-ranging studies are lacking.

While traditional use has established an empirical framework for dosage, variability in plant part used, harvesting season, preparation method, and individual constitution influences efficacy and tolerability. Hence, the emphasis in modern practice is shifting towards quality-controlled formulations, where dosage is defined not only by crude weight but also by active constituent content. Safety concerns, particularly related to tannin overconsumption, demand that decoctions and powders be used in moderation, with attention to duration of therapy.

8. Quality Control and Standardization

Ensuring the authenticity, purity, and consistency of *Helicteres isora* raw drugs is essential, given the plant's widespread ethnomedicinal use and growing demand in the herbal market. Quality control begins with proper botanical identification. The unique spirally twisted fruits serve as a macroscopic diagnostic feature, while microscopic characters—such as the presence of sclerenchyma fibers, calcium oxalate crystals, and tannin-filled parenchyma—aid in differentiating the crude drug from adulterants.

Pharmacopoeial standards provide baseline parameters for fruit quality. The *Ayurvedic Pharmacopoeia of India* specifies limits such as loss on drying not more than 12%, total ash below 10%, and alcohol-soluble extractives not less than 8%. Tannin content is typically in the range of 15-20% in fruits, which serves as an important quality marker.

Chemical standardization relies on chromatographic fingerprinting. High-Performance Thin-Layer Chromatography (HPTLC) and High-Performance Liquid Chromatography (HPLC) have been employed to develop reliable fingerprints using gallic acid, catechin, epicatechin, and betulinic acid as marker compounds. Recent studies using UPLC and LC-MS/MS have expanded the phytochemical profile to include multiple flavonoids and phenolic acids, which can be used for establishing more comprehensive quality benchmarks.

Standardization also requires careful monitoring of contaminants. Heavy metals, pesticide residues, microbial load, and aflatoxins should be within the limits prescribed by pharmacopoeial guidelines. Given the plant's popularity in rural markets, where adulteration with morphologically similar fruits or other astringent botanicals is common,

layered authentication approaches combining macroscopic, microscopic, and chromatographic markers are recommended. A summary of key quality control parameters is presented below.

Table 4: Quality Control and Standardization Parameters of *Helicteres isora*

Parameter	Specification / Marker	Reference Source
Loss on drying	Not more than 12%	Ayurvedic Pharmacopoeia of India (2007)
Total ash	Not more than 10%	API
Alcohol-soluble extractives	Not less than 8%	API
Tannin content	15-20% (fruits)	Gupta *et al*., 2013
Marker compounds	Gallic acid, catechin, epicatechin, betulinic acid	HPTLC/HPLC standards
Microscopy	Sclerenchyma fibers, calcium oxalate crystals, tannin-filled parenchyma	Pharmacognostic studies
Contaminants	Heavy metals, pesticides, aflatoxins, microbial load within pharmacopeial limits	WHO & API standards

The adoption of such rigorous standards is essential for ensuring consistency in therapeutic outcomes, facilitating global trade, and safeguarding consumer safety.

9. Conservation, Ethics, and Sustainable Supply

The growing popularity of *Helicteres isora* in traditional medicine, coupled with rising demand in the commercial herbal sector, has placed increasing pressure on wild populations. Fruits and bark are harvested extensively in central India, particularly in Madhya Pradesh and Chhattisgarh, which supply a significant share of the annual trade estimated at 200-250 metric tons. Field surveys conducted by the Foundation for Revitalization of Local Health Traditions (FRLHT) have indicated a decline in wild populations in several regions due to unsustainable harvesting practices, habitat degradation, and lack of systematic cultivation.

Ethical sourcing and conservation strategies are therefore essential to safeguard the long-term availability of this species. Good Collection Practices (GCP) recommend non-destructive harvesting methods, such as collecting only mature fruits and avoiding excessive bark stripping, which can damage or kill trees. Rotational harvesting cycles should be introduced to allow regeneration of natural populations.

Cultivation has emerged as a sustainable alternative to wild harvesting. The Indian Council of Forestry Research and Education (ICFRE) has reported that *H. isora* can be successfully propagated through seeds and stem cuttings, though natural seed germination rates remain moderate at around 40-50%. Pre-treatment with hot water has been shown to improve germination efficiency. Incorporating *H. isora* into agroforestry systems or contract farming schemes can reduce dependence on forest resources while providing economic benefits to rural communities.

Ethical considerations also extend to benefit-sharing with indigenous communities who have safeguarded the knowledge of this plant's uses for generations. Community-based cultivation projects and equitable pricing structures can ensure that local collectors and traditional healers are not marginalized by the commercialization of herbal medicines. Furthermore, adherence to biodiversity conventions and fair-trade practices strengthens the sustainability framework.

In conclusion, the conservation and ethical utilization of *Helicteres isora* require a multi-pronged approach involving habitat protection, sustainable harvesting, cultivation initiatives, and community engagement. Without such measures, increasing market demand risks overexploitation, loss of biodiversity, and erosion of traditional knowledge. Sustainable supply chains, coupled with ethical benefit-sharing, will not only secure the future of *H. isora* but also enhance its credibility and acceptance in global herbal therapeutics.

10. Conclusion

Helicteres isora is a credible ethnomedicinal resource with strong astringent, antioxidant, antimicrobial, and metabolic-modulating signatures that rationalize its traditional use in gastrointestinal and related indications. The plant's diverse polyphenolic and triterpenoid profile offers multiple mechanistic entry points for modern phytotherapeutic development. Realization of its full potential will require rigorous standardization, toxicity clarification, and well-designed clinical evaluations coupled with sustainable sourcing and ethical supply chains. In the interim, *H. isora* can be responsibly incorporated into evidence-informed, quality-controlled polyherbal formulations, particularly for acute, self-limiting gastrointestinal complaints and as an adjunct in metabolic health—under professional guidance and with attention to safety and quality.

Acknowledgments: Not applicable.

Declarations

Funding: None.

Conflicts of interest: None.

Ethical approval: Not applicable (narrative review).

References

1. Ayurvedic Pharmacopoeia of India. Part I, Vol. 5. New Delhi: Government of India, Ministry of Health and Family Welfare, Department of AYUSH; 2007. p. 45-47.
2. Gupta PC, Sharma N, Rao CV. A review on ethnobotany, phytochemistry and pharmacology of *Helicteres isora* Linn. J Pharmacogn Phytochem. 2013;2(4):64-74.
3. Kumar S, Yadav DK, Prasad DN, Hemalatha S. Pharmacognostical and phytochemical evaluation of *Helicteres isora* L. fruits. Pharmacognosy Rev. 2010;4(7):108-113.
4. Foundation for Revitalization of Local Health Traditions (FRLHT). Medicinal Plants in Trade: *Helicteres isora*. Bangalore: FRLHT; 2015. p. 32-34.
5. National Medicinal Plant Board (NMPB). Annual report 2014-15. New Delhi: Ministry of AYUSH; 2015. p. 67-70.
6. Indian Council of Forestry Research and Education (ICFRE). Propagation studies on medicinal plants of central India. Dehradun: ICFRE Publication; 2016. p. 12-14.

7. Ahmed F, Urooj A. Antioxidant properties and antimicrobial activities of *Helicteres isora*. J Food Biochem. 2009;33(6):790-808.
8. Bhakuni DS, Jain S, Sharma VN. Flavonoids from the fruits of *Helicteres isora*. Phytochemistry. 1972;11(2):667-669.
9. Choudhary GP, Sharma V, Kumar R. Antidiarrheal activity of fruit extracts of *Helicteres isora* Linn. in experimental animals. Indian J Pharmacol. 2011;43(2):192-195.
10. Devi BP, Boominathan R, Mandal SC. Evaluation of antidiabetic potential of fruit extract of *Helicteres isora* Linn. in streptozotocin-induced diabetic rats. J Ethnopharmacol. 2008;118(2):210-214.
11. Kalyani GA, Rajani M, Padh H. HPTLC method for fingerprinting of *Helicteres isora* fruits. Indian Drugs. 2007;44(9):693-697.
12. Sharma M, Jain V, Katoch V. Hepatoprotective effect of *Helicteres isora* against CCl_4-induced toxicity in albino rats. Int J Pharm Sci Res. 2012;3(5):1506-1512.
13. Singh R, Chaurasiya ND, Sangwan NS, Tiwari M, Verma RK. Profiling of bioactive compounds in *Helicteres isora* fruits by LC-MS/MS. Planta Med. 2014;80(16):1475-1482.
14. World Health Organization (WHO). Quality control methods for medicinal plant materials. Geneva: WHO; 2011. p. 21-28.
15. Kadus PA, Giramkar AA, Nirmal SA. A review on: Avartani (*Helicteres isora*). Int J Agric Nutr. 2023;5(1):55-58. https://doi.org/10.33545/26646064.2023.v5.i1a.100
16. Venkatesh S, Reddy GD, Ramesh M, Apparao AVN. Antihyperglycemic activity of *Helicteres isora* root in alloxan-induced diabetic rats. Pharm Biol. 2003;41(8):628-631.